

# Pure

## Scotland's Rural College

### **A critical review of the impacts of cover crops on nitrogen leaching, net greenhouse gas balance and crop productivity**

Abdalla, M; Hastings, A; Cheng, Kun; Yue, Q; Chadwick, David; Espenberg, Mikk; Truu, Jaak; Rees, RM; Smith, Pete

*Published in:*  
Global Change Biology

*DOI:*  
[10.1111/gcb.14644](https://doi.org/10.1111/gcb.14644)

Print publication: 01/08/2019

*Document Version*  
Peer reviewed version

[Link to publication](#)

#### *Citation for pulished version (APA):*

Abdalla, M., Hastings, A., Cheng, K., Yue, Q., Chadwick, D., Espenberg, M., ... Smith, P. (2019). A critical review of the impacts of cover crops on nitrogen leaching, net greenhouse gas balance and crop productivity. *Global Change Biology*, 25(8), 2530-2543. <https://doi.org/10.1111/gcb.14644>

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# **A critical review of the impacts of cover crops on nitrogen leaching, net greenhouse gas balance and crop productivity**

Mohamed Abdalla<sup>a</sup>, Astley Hastings<sup>a</sup>, Kun Cheng<sup>b</sup>, Qian Yue<sup>b</sup>, Dave Chadwick<sup>c</sup>, Mikk Espenberg<sup>d</sup>, Jaak Truu<sup>d</sup>, Robert M Rees<sup>e</sup>, Pete Smith<sup>a</sup>

<sup>a</sup>Institute of Biological and Environmental Sciences, School of Biological Sciences, University of Aberdeen, Aberdeen, AB24 3UU, UK.

<sup>b</sup>Institute of Resource, Ecosystem and Environment of Agriculture, and Centre of Climate Change and Agriculture, Nanjing Agricultural University, Nanjing, Jiangsu, China.

<sup>c</sup>School of Natural Resources, Bangor University, Bangor, Gwynedd, LL57 2UW, UK.

<sup>d</sup>Faculty of Science and Technology, University of Tartu, Tartu, Estonia.

<sup>e</sup>Scotland's Rural College (SRUC) Edinburgh, West Mains Road, Edinburgh, EH93JG, UK.

**Key words:** Cover crop, Catch crop, N leaching, Soil organic carbon, N in grain, Nitrous oxide emissions, net greenhouse gas balance, green manure, yield, N content, nitrate, C sequestration

## **Abstract**

Cover crops play an increasingly important role in improving soil quality, reducing agricultural inputs and improving environmental sustainability. The main objectives of this critical global review and systematic analysis were to assess cover crop practices in the context of their impacts on nitrogen leaching, net greenhouse gas balances (NGHGB) and crop productivity. Only studies that investigated the impacts of cover crops and measured one or a combination of: nitrogen leaching, soil organic carbon (SOC), nitrous oxide (N<sub>2</sub>O), grain yield and nitrogen in grain of primary crop, and had a control treatment were included in the analysis. Long-term studies were uncommon, with most data coming from studies lasting 2-3 years. The literature search resulted in 106 studies carried out at 372 sites and covering different countries, climatic

zones and management. Our analysis demonstrates that cover crops significantly ( $p < 0.001$ ) decreased N leaching and significantly ( $p < 0.001$ ) increased SOC sequestration without having significant ( $p > 0.05$ ) effects on direct  $\text{N}_2\text{O}$  emissions. Cover crops could mitigate the NGHGB by  $2.06 \pm 2.10 \text{ Mg CO}_2\text{-eq ha}^{-1} \text{ y}^{-1}$  and significantly ( $p < 0.05$ ) increase N in grain of the primary crops. One of the potential disadvantages of cover crops identified was the reduction in grain yield of the primary crop by  $\approx 3.9\%$ , compared to the control treatment. This drawback could be avoided by selecting mixed cover crops with a range of legumes and non-legumes, which increased the yield by  $\approx 13\%$ . These advantages of cover crops justify their widespread adoption. However, management practices in relation to cover crops will need to be adapted to specific soil, management and regional climatic conditions.

## **Introduction**

Increasing crop productivity with reduced inputs and lower impacts on the environment is a major current challenge for global food production. Cover crops (also known as catch crops) are plants mostly grown after a primary crop is harvested, in regions of the world where only a single main crop is grown (such as North Europe, North China and Canada). This avoids periods of bare soil which are associated with greater risk of erosion and nitrogen leaching losses (Battany and Grismer, 2000). Cover cropping can comprise of a single species or a mixture of species and can use annual, biennial, or perennial vegetation. Cover crops can be killed (or ploughed-in) in winter or spring, or grazed, and incorporated in soils by tillage to prevent competition with the primary crop, and to promote mineralization of organic N (Dabney et al., 2011). They can also left on the soil surface over the fall and winter periods, until a primary crop in no-till is planted, to provide weed control and N inputs (Halde et al., 2014).

55

56 Cover crops can increase water holding capacity, soil porosity, aggregate stability, the size of  
57 the microbial population and its activity and nutrient cycling (Lotter et al., 2003; Drinkwater  
58 and Snapp, 2007; Harunaa & Nkongolo, 2015). There are four classes of cover crops: legumes  
59 (e.g. alfalfa, vetches and clover), non-legumes (spinach, canola and flax), grasses (e.g. ryegrass  
60 and barley) and brassicas (e.g. radishes and turnips). The two main types of cover crops are  
61 legumes and non-legumes. Legume cover crops have the ability to fix nitrogen (N) biologically  
62 and increase soil organic matter (SOM) content (Lüscher et al., 2014). They can be used as a  
63 green manure to improve soil nutrition for the subsequent primary crop. On the other hand,  
64 non-legume cover crops can absorb excess nitrate from the soil, increase crop biomass, and  
65 improve soil quality (Finney et al., 2016; White et al., 2016). Farmers, generally, select specific  
66 types of cover crops based on their own needs and goals influenced by biological,  
67 environmental, social, cultural and economic factors of the farming systems in which they  
68 operate (Snapp et al., 2005). Additionally, cover crops have become of greater interest for their  
69 potential to provide additional ecosystem services in agricultural systems (e.g. reduce erosion,  
70 improve water quality and enhance biodiversity). In Spain, Hontoria et al. (2019) found that  
71 the use of barley as a winter CC is an appropriate choice to promote arbuscular mycorrhizal  
72 fungal populations and biological activity in soils with intercropping systems.

73

74 Nitrogen leaching from agricultural soils is of great concern due to its contribution to  
75 excess nitrate ( $\text{NO}_3$ ) concentrations in ground water and run-off (Ascott et al., 2017), indirect  
76 emissions of greenhouse gases (GHGs) e.g. nitrous oxide ( $\text{N}_2\text{O}$ ) (Delgado et al., 2008), and  
77 loss of expensive N fertilizer (Cardenas et al., 2011). This problem is more pronounced in areas  
78 with fertilized coarse-textured soils (Basche et al., 2014) or areas with high precipitation  
79 (Thorup-Kristensen et al., 2003). In England, Allingham et al. (2002) reported an average  $\text{NO}_3$

leaching value of 65 kg N ha<sup>-1</sup>, which is approximately 25% of total N input. Similar NO<sub>3</sub> losses, as a proportion of the total N applied, have been reported following livestock slurry and poultry manure applications to arable soils (Chambers et al., 2000). Previous studies have found that replacing fallow periods with non-legume cover crops is an effective management practice to withdraw soil N into the biomass of the cover crops and to reduce NO<sub>3</sub> leaching (Kaspar and Singer, 2011; Quemada et al., 2013; Basche et al., 2014). Cover crops can also increase soil organic carbon (SOC) stocks in agricultural soils (Poeplau and Don, 2015), since more C and N are added to the soil pools as cover crop residues decompose (Steenwerth and Belina, 2008; Kaspar and Singer, 2011). The amounts of C and N incorporated into the soil depend on many factors e.g. the amount, quality and management of the residues, soil type, frequency of tillage and climatic conditions (Stevenson, 1982; Smith et al., 1996). However, it is still not clear how cover crops affect the net greenhouse gas balance (NGHGB). Further, there is conflicting evidence on the influence of the cover crops on grain yields and N in the grain of primary crops. Some previous studies found that under-sowing of cover crops in spring could lead to a high level of competition with the primary crop for nutrients, soil moisture and light, and result in some loss of the grain yield (Karlsson-Strese et al., 1998; Känkänen et al., 2001, 2003). Other studies found that grain yield of the primary crops was not affected (Wallgren and Lindén, 1994; Ohlander et al., 1996) or was even increased (Campiglia et al., 2011). Mixed results have also been reported for the effects of cover crops on N in grain of the primary crop (Thomsen, 2005; Rinnofner et al., 2008; Doltra and Olesen, 2013).

The main objectives of this global review and systematic analysis were to investigate the impacts of cover crops (legume, non-legume and legume-non-legume mixed) on N leaching, the NGHGB and crop productivity in terms of grain yield and N content in the grain of the primary crop. We also investigated whether soil characteristics, field management and

climatic zones can modify these effects, and through this, we assessed the viability of cover crops as a management tool to enhance C sequestration, reduce N loss from agroecosystems and maintain crop production. The specific hypotheses we critically evaluated were as follows: 1) cover crops decrease N loss and increase SOC accumulation, 2) the impacts of cover crops on N loss and SOC are modified by soil, management and climatic zones, and 3) including cover crops in crop rotations improves grain yield and N in grain of the primary crop.

## **Materials and Methods**

### *Data collection*

To analyse the publications that have investigated the impacts of cover crops on N leaching, SOC, N<sub>2</sub>O, grain yield and N in grain for different primary crops (e.g. wheat, barley, oats, corn and others), we made a comprehensive search on the Web of Science database (accessed between January, 2017 and September, 2018) using the keywords: Cover crop, Catch crop, N leaching, Soil organic carbon, N in grain, Nitrous oxide emissions, GHG balance, green manure, yield, N content, nitrate and C sequestration. To gain the best possible coverage of the topic, we also checked all references in the papers collected from the Web of Science search. We only selected studies that investigated the effects of cover crops (legume, non-legume and legume-non-legume mixed), covered at least one growing season and measured one or a combination of: N leaching, SOC, N<sub>2</sub>O, grain yield and N in grain of primary crop, and had a control treatment. Nitrous oxide data were collected from studies that measured the flux from cropland and applied either a static or automated chamber method. In some studies SOC values are given as concentrations. To convert these values to stocks (t ha<sup>-1</sup>), we applied equation 1 below (Guo & Gifford, 2002):

$$Cs = (SOC * BD * D)/10 \quad (1)$$

Where, Cs is soil organic carbon stocks ( $\text{Mg ha}^{-1}$ ); SOC is soil organic carbon concentration ( $\text{g kg}^{-1}$ ); BD ( $\text{g cm}^{-3}$ ); and D is soil depth (cm).

For SOC and N leaching data, we selected studies that measured them from zero and up to 30 and 100 cm soil depth, respectively. To improve comparability of the different studies, we normalized the SOC data to the top 30 cm and the N leaching data to the top 100 cm depth, using the depth distribution method produced by Jobbágy and Jackson (2001) (equations 2-4).

$$Y = 1 - \beta^d \quad (2)$$

$$\text{SOC}_{30} = ((1 - \beta^{30}) / (1 - \beta^{d_0})) * \text{SOC}_{d_0} \quad (3)$$

$$\text{N}_{100} = ((1 - \beta^{100}) / (1 - \beta^{d_0})) * \text{N}_{d_0} \quad (4)$$

Where Y is the cumulative proportion of the SOC or soil N leaching pool from the soil surface to depth d (cm);  $\beta$  is the relative rate of decrease in the soil SOC or N pool with soil depth (0.9786 for SOC and 0.9831 for N) (Jobbágy and Jackson 2000; Jobbágy and Jackson 2001).  $\text{SOC}_{30}$  or  $\text{N}_{100}$  is the SOC ( $\text{t ha}^{-1}$ ) or N ( $\text{kg N ha}^{-1}$ ) pool in the upper 30 or 100 cm depth, respectively;  $d_0$  is the original soil depth available in individual studies (cm);  $\text{SOC}_{d_0}$  or  $\text{N}_{d_0}$  is the original soil SOC or N pool.

We defined the control treatment as an annual fertilized primary crop with a bare fallow period between harvest and the establishment of the next primary crop. Where two main crops are grown synchronously, they are usually then referred to as intercrops, and such systems were not considered further in this review. We excluded many studies either because there was no control or because the experimental treatments did not meet the above criteria. Our literature search resulted in 106 studies carried out at 372 sites (Tables S1-S5) that investigated the

impacts of cover crops on N leaching, grain yield and N in grain of primary crop, SOC, N<sub>2</sub>O emissions, respectively, and covering different countries, climatic zones and management systems. The majority of the studies collected were short-term experiments of 2-3 years. Locations, climatic conditions as well as primary crop, cover crops, type of cover crops (legume, non-legume or legume-non-legume mixed), study duration, tillage, N fertilizer application rate, soil texture, soil depth (cm), bulk density (BD), soil pH and measurements from control and treatments i.e. N leaching, grain yield, N in grain of primary crop, SOC and N<sub>2</sub>O, are shown in Tables S1-S5. When there was more than one year of study in the original paper, we used the mean value for different years. We included different methods for measuring N leaching (e.g. field cores, ceramic suction cup lysimeter, and subsurface drainage lysimeter). Nitrogen leaching was measured/ calculated in kg N ha<sup>-1</sup> y<sup>-1</sup> whilst SOC and grain yield in t ha<sup>-1</sup> y<sup>-1</sup>, and N in grain in g N m<sup>-2</sup> y<sup>-1</sup>. We found 78% of the N leaching dataset collected had conventional tillage systems whilst the rest (22%) was divided between the different types of conservation tillage systems (i.e. no-till, reduced till and minimum till) or had no data. Therefore, we investigated the influence of tillage on cover crop efficiency to reduce N leaching, N<sub>2</sub>O and SOC by comparing between conventional and conservation tillage systems.

To investigate the impacts of climate, we divided our dataset into four groups depending on the climatic zones. Climatic zones were distinguished on the basis of temperature and moisture regimes (cool, warm, dry and moist zone) to represent the global variations of soil moisture and temperature. The cool zone covers the temperate (oceanic, sub-continental, and continental) and boreal (oceanic, sub-continental and continental) areas, whilst the warm zone covers the tropics (lowland and highland) and subtropical (summer rainfall, winter rainfall, and low rainfall) areas (Smith et al., 2008; Abdalla et al., 2018). The dry zone includes



the areas where the annual precipitation is  $\leq 500$  mm, whilst the moist zone includes areas where the annual precipitation is  $> 500$  mm (Smith et al., 2008). The four climate categories were; moist cool (MC), moist warm (MW), dry cool (DC) and dry warm (DW). However, to investigate the influences of climatic zones on the efficiency of cover crops to reduce N leaching and SOC, comparisons were made between the MC and MW only as most of the dataset belong to these two climatic zones: MC (68%) and MW (24%). The two other climatic zones both have only four observations.

For the different studies, different methods were used to measure soil pH e.g. using a pH probe or meter in deionized water or 0.01 M  $\text{CaCl}_2$  in 1:1 and 1:2 or 1:5 (v:v) soils: solution ratios. We assumed the pH results to be equivalent, and where a range of values were reported we took the arithmetic mean. The mean annual air temperature (MAAT, in  $^{\circ}\text{C}$ ) value, and mean annual precipitation (MAP, in mm) values for each study, were collected from the original published papers. The locations of experiments used in this study were plotted on a map of net primary production (NPP) calculated using the Miami method (Lieth, 1972; Grieser et al., 2006), to indicate the diversity of arable capability included (Fig. 1).

## *2.2 Direct/ indirect $\text{N}_2\text{O}$ emissions and net greenhouse gas balance (NGHGB)*

The direct  $\text{N}_2\text{O}$  emissions data were collected from the literature (Table S5). Following Tier I IPCC protocol (IPCC, 2006) and Parkin et al. (2016), we estimated the indirect  $\text{N}_2\text{O}$  emissions for the control and cover crop treatments from the N leaching using the EF of 0.0075 multiplied by the mass of N leached. The change in the indirect  $\text{N}_2\text{O}$  emissions due to cover crops were then calculated as shown in Table S1. The indirect emissions associated with  $\text{NH}_3$  and  $\text{NO}_x$  were not estimated. The contributions of SOC (Table S4) and  $\text{N}_2\text{O}$  to the NGHGB were calculated using the IPCC (2013) approach, where on a mass basis,  $\text{N}_2\text{O}$  has a global warming

potential (GWP) of 298 times that of CO<sub>2</sub>, over a 100-year timescale. The methane (CH<sub>4</sub>) flux was considered to be negligible as, generally, cropland soils tend to be well drained and oxygenated and are often small net CH<sub>4</sub> sinks (Lee et al., 2006; Abdalla et al., 2014). The NGHGB was calculated as the difference between the increases in GWP due to higher direct N<sub>2</sub>O emissions and the decreases due to higher SOC accumulation and lower indirect N<sub>2</sub>O emissions under the cover crops.

### *Data analyses*

We used R version 3.5.2 (R Development Core Team, 2018) to perform exploration, harmonisation and analyses of the data. The distributions of N leaching, grain yield, N in grain, N<sub>2</sub>O and SOC measurements were characterised using the “fitdistrplus” package version 1.0-14 (Delignette-Muller and Dutang 2015). To investigate difference on all sites where both the control and cover crop treatments (cover crop types, climatic zones, tillage systems) had N leaching, grain yield, N in grain, N<sub>2</sub>O and SOC measurements, we used the “glmer” method with random effect (different studies) and Gamma (link “log”) distribution (version 1.1-19) (Bates et al., 2015), while p-values were calculated in order to confirm the significance of the relationships using the “lmerTest” package version 3.0-1 (Kuznetsova et al., 2017). The same method was performed to test whether there was a significant difference in N leaching, grain yield, N in grain, N<sub>2</sub>O emissions and SOC between cover crops, tillage, climatic zones and soil texture types. A linear mixed effects model function was applied to investigate whether there was an effect of cover crops, tillage, climatic zones and soil texture types on physicochemical values. Analysis of covariance (ANCOVA) was used to compare N leaching (%) of cover crops (legume, non-legume and legume-non-legume mixed), with added N fertilizer as covariate in the model. The package “akima” version 0.6-2 was used to create interpolated contour plots (Akima and Gebhardt, 2015) of pairs of the BD, pH and added N as x-axis and y-axis with N

leaching and SOC as the z variable. A contour plot is a graphical technique for representing a 3-dimensional surface by plotting constant z slices on a 2-dimensional format. That is, given a value for z, lines are drawn for connecting the (x,y) coordinates where that z value occurs. We performed linear regressions of different variables against N leaching and SOC.

## Results


### *Impacts of cover crops (legume, non-legume and legume-non-legume mixed) on N leaching*

The inclusion of cover crops in the crop rotation significantly decreased N leaching compared to the control treatments ( $p < 0.001$ ;  $n = 75$ ). All types of cover crops had significant effects on N leaching; legume ( $p < 0.05$ ;  $n = 11$ ), non-legume ( $p < 0.001$ ;  $n = 55$ ) and legume-non-legume mixed cover crops ( $p < 0.001$ ;  $n = 9$ ) (Fig. 2a). A one-way ANOVA showed no significant ( $p > 0.05$ ) difference in N leaching between legume, non-legume and legume-non-legume mixed cover crops. Additionally, an analysis of covariance (ANCOVA) showed no significant ( $p > 0.05$ ) effect of cover crops on the change of N leaching (%), after controlling for the effect of added N fertilizer application rate (the covariate) ( $F = 1.23$ ,  $p = 0.3$ ) (Fig. 3).


### *Impacts of cover crops (legume, non-legume and legume-non-legume mixed) on SOC and direct N<sub>2</sub>O emissions*

A paired t-test showed that SOC under the cover crops was significantly higher compared to that in the control treatments ( $p < 0.001$ ;  $n = 43$ ). Both legume ( $p < 0.001$ ,  $n = 29$ ) and non-legume ( $p < 0.001$ ;  $n = 13$ ) cover crops significantly increased SOC (Fig. 2d). A paired t-test showed that cover crops ( $n = 28$ ) had no significant effect ( $p > 0.05$ ) on direct N<sub>2</sub>O emissions, compared to the control treatment. Only legume ( $n = 8$ ) cover crops significantly increased direct N<sub>2</sub>O

254 emissions but non-legume (n=17) and legume-non-legume had no effects, compared to the  
255 control treatment.




**Fig. 1:** Map showing the net primary productivity (NPP) and locations of experimental sites considered in this paper. NPP calculated using the Miami method (Lieth, 1972; Grieser et al., 2006).



**Fig. 2:** Comparisons between N leaching (a), grain yield (b), N in grain (c) and SOC (d) from control and cover crops (CC) treatments. Types of cover crops (legume (blue), non-legume (green) or mixed (red)) and their 95% confidence intervals (CI).

Tillage had no effect on direct N<sub>2</sub>O emissions. However, the changes in direct N<sub>2</sub>O emissions (%) under conservation tillage were significantly lower compared to that under conventional tillage treatment (Table 1).



**Fig. 3:** Relationships between change in N leaching (%) and legume, non-legume and mixed cover crops. Least squares means of N leaching after analysis of covariance (ANCOVA,  $F=1.23$ ,  $n=86$ ,  $p=0.3$ ) with added N fertilizer used as covariates (vertical bars denote 95% confidence intervals).

Table 1: Effects of tillage on direct  $N_2O$  emission ( $kg\ ha^{-1}y^{-1}$ ) from control and cover crop treatments.

Treatment	Mean $\pm$ StDev. (conventional)	N* (conventional)	Mean $\pm$ StDev. (conservation)	N (conservation)	t-value	p
Control	0.94 $\pm$ 1.0	12	3.70 $\pm$ 2.74	10	3.25	ns
Cover crops	1.46 $\pm$ 1.61	12	3.95 $\pm$ 2.91	10	2.55	ns
Change in $N_2O$ emissions (%)	50.58 $\pm$ 148.34	12	16.65 $\pm$ 38.94	10	4.74	$p<0.001$

N\* = number of observation; StDev. = standard deviation; ns= not significant.

*Impacts of cover crops (legume, non-legume and legume-non-legume mixed) on grain yields and N in grain of primary crop*


Overall, the cover crops significantly decreased grain yield of the primary crops compared to the control treatments (on average -3.9%;  $p < 0.05$ ;  $n = 154$ ) (Fig. 2b). Both legume and non-legume cover crops significantly decreased ( $p < 0.001$ ;  $n = 52$  and  $p < 0.01$ ;  $n = 96$ , respectively) grain yield of the primary crop whilst legume-non-legume mixed cover crops significantly increased ( $p < 0.01$ ,  $n = 6$ ) grain yield of the primary crop (by  $\approx 13\%$ ). Cover crops significantly ( $n = 118$ ;  $p < 0.001$ ) decreased grain yield of the primary crop under conventional tillage but had no effect under conservation tillage ( $n = 20$ ;  $p > 0.05$ ). The cover crops, generally, had no effect on N content in the grain of the primary crop (n.s;  $n = 58$ ) (Fig. 2c). Though, both legume and non-legume cover crops significantly increased N in the grain of the primary crop ( $p < 0.001$ ;  $n = 15$  and  $p < 0.05$ ;  $n = 39$ , respectively). Legume-non-legume mixed cover crops had no effects ( $p > 0.05$ ;  $n = 4$ ) on N in grain of the primary crop.

*Influences of management, soil and climatic zones on cover crop efficiency to decrease N leaching and to increase SOC*


For N leaching at 0-100 cm depth, contour plots based on available data, showed that BD and N fertilizer application rate explained 11.6% of overall variance ( $p < 0.01$ ;  $n = 38$ ). N leaching was significantly related to BD ( $p < 0.05$ ) (Fig. 4). For the SOC at 0-30 cm depth, BD and N fertilizer application rate explained 57% of the overall variance in SOC ( $p < 0.001$ ;  $n = 41$ ). The increase in SOC under cover crops was significantly related to both N fertilizer application rate ( $p < 0.01$ ) and BD ( $p < 0.001$ ) (Fig. 5). The interaction between soil pH and N fertilizer application rate had no significant effect on N leaching ( $p > 0.05$ ;  $n = 43$ ). Soil pH and added N fertilizer application rate significantly influenced SOC and explained 31% of the overall variance ( $p < 0.01$ ;  $n = 35$ ). However, changes in SOC varied significantly with soil pH ( $p < 0.001$ )




(Fig. 6). Soil texture had no significant ( $p>0.05$ ) impacts on the change in N leaching or SOC. The N leaching and SOC under the control and cover crop treatments were both not significantly ( $p>0.05$ ) influenced by MAAT.



**Fig. 4:** Contour plot ( $n=38$ ) showing relationships between added N fertilizer application rate, BD and change in N leaching (%) at 0-100 cm depth. These two variables explain 11.6% of N leaching overall variation ( $p<0.05$ ). N leaching significantly depended on BD ( $t=2.62$ ;  $p<0.01$ ). One outlier was removed ( $BD=2.5$ ).



**Fig. 5:** Contour plot (n=41) showing relationships between added N fertilizer application rate, BD and change in SOC (%). Added N fertilizer and BD explain 57% of SOC overall variation ( $p < 0.001$ ). The SOC depended significantly on added N ( $t = -3.2$ ;  $p < 0.01$ ) and BD ( $t = 7.1$ ;  $p < 0.001$ ).




**Fig. 6:** Contour plot (n=35) showing relationships between added N fertilizer application rate, pH and change in SOC. Added N fertilizer and pH explain 31% of SOC overall variation. SOC depended significantly on pH ( $t = -3.94$ ;  $p < 0.001$ ).

Cover crops significantly decreased N leaching under both MW ( $p < 0.001$ ;  $n = 13$ ) and MC ( $p < 0.001$ ;  $n = 58$ ) climatic zones. MAP positively correlated with SOC for the control ( $r^2 = 0.39$ ,  $p < 0.001$ ;  $n = 43$ ), and cover crops ( $r^2 = 0.39$ ,  $p < 0.001$ ;  $n = 43$ ) treatments (Fig. 7). Cover crops significantly increased SOC under MW ( $p < 0.001$ ;  $n = 37$ ) and under MC ( $p < 0.001$ ;  $n = 6$ ) climatic zones. Under both the conventional ( $n = 62$ ) and conservation ( $n = 12$ ) tillage systems, cover crops significantly ( $p < 0.001$ ) decreased N leaching compared to the control. A t-test showed that conservation tillage ( $n = 62$ ) significantly increased N leaching for the control ( $p < 0.05$ ) treatment compared to conventional tillage ( $n = 12$ ). There were no significant ( $p > 0.05$ ) effects on SOC due to tillage systems. The SOC was significantly higher under both

the conventional ( $p<0.05$ ,  $n=18$ ) and conservation ( $p<0.01$ ,  $n=17$ ) tillage systems compared to the control.

#### *Impacts of cover crops on net greenhouse gas balance*

Cover crops increased SOC and decreased N leaching and thereby, lowered the indirect N<sub>2</sub>O emissions (i.e. from N leaching) without significantly increasing direct N<sub>2</sub>O emissions. This combination of higher SOC and the lower indirect N<sub>2</sub>O emissions under the cover crops resulted in a lower NGHGB compared to the control treatment. The estimated reduction in NGHGB due to cover crops, compared to the control treatments, was  $2.06 \pm 2.10$  Mg CO<sub>2</sub>-eq ha<sup>-1</sup> y<sup>-1</sup>. The reduction in NGHGB due to different cover crop types, compared to the control treatments, were  $1.87 \pm 1.82$ ,  $1.82 \pm 1.44$  and  $5.15 \pm 3.51$  Mg CO<sub>2</sub>-eq ha<sup>-1</sup> y<sup>-1</sup> for the legume, non-legume and legume-non-legume mixed cover crops, respectively (Table 2). No significant difference ( $p>0.05$ ) was found between the different cover crop types.



**Fig. 7:** Relationships between SOC and mean annual air temperature (MAAT) (a) and mean annual precipitation (MAP) (b) under control and cover crops. MAAT was not significantly correlated with SOC ( $p>0.05$ ). MAP was positively correlated with SOC for both the control ( $t=5.0$ ,  $p<0.001$ ;  $r^2=0.39$ ,  $p<0.001$ ,  $n=43$ ), and cover crop ( $t=5.0$ ,  $p<0.001$ ;  $r^2=0.39$ ,  $p<0.001$ ,  $n=43$ ).

Table 2: Descriptive statistics of the reduction in net greenhouse gas balance (NGHGB) related to the reduction of indirect nitrous oxide ( $N_2O$ ) emission and the soil organic carbon sequestration ( $Mg\ CO_2\text{-eq}\ ha^{-1}\ y^{-1}$ ).

Type of cover crop	Change in direct N <sub>2</sub> O (mean±StDev*)	Change in indirect N <sub>2</sub> O (mean±StDev*)	Change in SOC (mean±StDev*)	N**	NGHGB (mean±StDev*)
Legume	0.04±0.05	-0.30±0.37	1.61±1.82	30	1.87±1.82
Non-legume	0.09±0.11	-0.07±0.28	5.12±5.51	13	1.82±1.44
Mixed	0.04±0.03	-0.50±0.37	0.30±0.37	4	5.15±3.51
All types	0.08±0.10	-0.16±0.33	1.97±2.10	47	2.06±2.10

\*StDev. = standard deviation. Negative numbers represent gas emissions, while positive numbers represent gain of C by the soil. \*\*N is the number of observations.

## Discussion

### *Impacts of cover crops (legume, non-legume and legume-non-legume mixed) on N leaching*

In this critical global review and systematic analysis, we found that all types of cover crops significantly decreased N leaching. However, no statistically significant differences between legume, non-legume and legume-non-legume mixed cover crops were found. Previous studies reported that non-legume (Torstensson and Aronsson, 2000; Aronsson et al., 2011; Thomsen and Hansen, 2014) legume (Salmerón et al., 2010; Askegaard and Eriksen, 2008; Askegaard et al., 2005) and legume-non-legume mixed (Askegaard et al., 2011; Benoit et al., 2014) cover crops can all reduce N leaching, but with different efficiencies. In the USA, Kaspar et al. (2012) reported that the use of non-legume cover crops (e.g. oat and rye) is a suitable management option for reducing N leaching from corn-soybean rotations and thereby, improving both water and soil quality. Non legume cover crops reduced soil NO<sub>3</sub> content which is vulnerable to N leaching during autumn and winter (Thorup-Kristensen et al., 2003), and made additional soil N available for the primary crop following mineralisation of their residues (Kaspar and Singer, 2011). In studying future scenarios over a period of 45-years, Tribouillois et al. (2018) found that non-legume cover crops continuously decreased N leaching compared to that of bare soil, but legume cover crop scenarios did not. Moreover, some simulation studies have suggested

that the efficiency of legume cover crops species to reduce N leaching was about half of that of non-legume species (e.g. Brassicaceae and Poaceae; Justes et al., 2012). Nevertheless, Valkama et al. (2015) reported that legume cover crops may not be effective in reducing N leaching but growing non-legume cover crops within a spring cereal crop is an effective method for reducing N leaching from different crop varieties, soils and weather conditions. Here, it is accepted that there is a trade-off between potential grain yield loss and environmental benefits, but this could be compensated for in environmental stewardship schemes in those countries. Leslie et al. (2017) recommended growing cover crops in some years only, to avoid a pre-emptive competition where the cover crops could recover soil NO<sub>3</sub> that would otherwise have been available to the subsequent primary crop. The non-legume cover crops can also increase N leaching when grown too late in spring or in dry areas, where the risk for N leaching is low (Thorup-Kristensen, 2003). Thus, the timing and location of the non-legume cover crops need to be considered carefully to avoid competition with the primary crop.

#### *Impacts of cover crops (legume, non-legume and legume-non-legume mixed) on SOC and direct N<sub>2</sub>O emissions*

Cover crops (i.e. both legume and non-legume) increased SOC, and so they can enhance C sequestration in soils. Similar conclusions regarding the impact of cover crops on SOC were reported by Wortman et al. (2012), Olson et al. (2014), Poeplau and Don (2015) and others. According to Ding et al. (2006), both organic carbon and light fraction C contents increased in soils under cover crops, with or without N fertilizer. Here, the decomposition of dead roots and biomass of cover crops result in improved SOM quantity and quality (Villamil et al., 2006). This could help improve food security, reduce NGHGB and mitigate climate change.

We found that cover crops had no significant effect on direct N<sub>2</sub>O emissions compared to the control. According to Webb et al. (2000), cover crops increase the direct N<sub>2</sub>O emissions when residues are incorporated into the soil or by increasing the photo-synthetically-derived C supply from actively growing root systems. However, adjusting the N fertilizer application rate (e.g. by integrated soil fertility management) could help in reducing the gas emissions (Guardia et al., 2016; Tribouillois et al., 2018). Previous studies reported contrasting results with regard to cover crop effects on direct N<sub>2</sub>O emissions (Abdalla et al., 2013; Mitchell et al., 2013; Basche et al., 2014). This could be explained by the large variations in many factors, e.g. cover crop types and performances, climate, soil characteristics, tillage and seasons of N<sub>2</sub>O samplings, between the different studies. Cover crops have the ability to decrease the indirect N<sub>2</sub>O emissions (i.e. from N leaching). Cover crop species influence abiotic and biotic soil factors differently (Abalos et al., 2014). They have the capacity to simultaneously mitigate N leaching and indirect N<sub>2</sub>O emissions (Kim et al., 2015) by limiting N availability. They deplete the soil NO<sub>3</sub> pool which is the major substrate for denitrification (Liebig et al., 2015), reducing N leaching and consequently decreasing the contribution of indirect N<sub>2</sub>O emissions to the NGHGB. However, this depends on many factors, e.g. cover crop types, performances, climate, tillage and soil characteristics. In contrast, Zhou and Butterbach-Bahl (2013) found that for coarser textured soils, the reduction in N leaching can increase availability of soil N, which can lead to a trade-off by enhancing N<sub>2</sub>O emissions.

#### *Influences of management, soil and climatic zones on cover crop efficiency to decrease N leaching and increase SOC*

Cover crops were most efficient in reducing N leaching when the BD was <1.4 g cm<sup>-3</sup> and N fertilizer application rate was >200 kg N ha<sup>-1</sup>. Snapp (2005) found application of more N fertilizer, especially with legume cover crops, can increase the risk of nutrient leaching, if a



subsequent primary crop is not planted promptly. Thus, to reduce N leaching from soils under cover crops, judicious quantities of N fertilizer should be applied at appropriate application times, with appropriate methods (Yogesh and Juo, 1982; Fan et al., 2010). Also, to avoid losing the excess N in soils by leaching, the amount of N fertilizer applied should be based on soil and crop requirement tests (Bundy et al., 2005; Defra, 2010).

In this study, we found enough data points for MW and MC climatic zones but not for DW and DC climatic zones. This is obviously because cover crops are rarely grown in dry climates as they use water that could be used to grow a primary crop and reduce water percolation by transpiration (Weinert et al., 2002). Additionally, in such climates, cover crops compete with the primary crop for nutrients (Unger and Vigil, 1998) and consequently, have negative impacts on crop growth and productivity. Wortman et al. (2012) and Tribouillois et al. (2018) reported that the large quantity of soil water used by the cover crops, at the cost of the subsequent primary crop and immobilisation of soil N due to incorporation of low quality cover crop residues into the soil is also a major concern. These problems appear mostly in arid and semiarid environments (< 500 mm annual rainfall) where water storage in soils declines with the establishment of cover crops, and results in reduced crop yields (Cherr et al., 2006; Nielsen and Vigil, 2005). Conservation tillage significantly decreased the efficiency of cover crops to decrease N leaching under control treatment compared to that under conventional tillage. The large pores that can develop under conservation tillage result in high N leaching if present after broadcasting N fertilizer (CTS, 2011), and thereby could also increase GHG emissions (Smeaton et al., 2011). Fraser et al. (2013) found that tillage had some effects on N leaching, though the use of minimum tillage for autumn cultivation resulted in significantly less N leaching than either intensive or no-till. Buchi et al. (2018) reported that cover crop could maintain wheat yield and improve soil fertility and nutrient cycling in a no-till system.

Therefore, a combination of the right type of conservation tillage with cover crops could be the best management to reduce N leaching in dry climates. Water utilization by the cover crops is counterbalanced by the improved infiltration and reduced evaporative losses that occur in conservation tillage systems (Unger and Vigil, 1998; Wang and Ngouajio, 2008). Further, the high soil moisture under conservation tillage positively influences microbial activity (Madejon et al., 2009) and increase bypass flow (CTS, 2011). This could also slow the rate of mineralization, as soils take longer to warm in the spring (Abdalla et al., 2013).

We found no significant effects on the efficiency of cover crops to decrease N leaching between the MW and MC climate zones. Fraser et al. (2013) and Hooker et al. (2008) found that inter-annual weather variability and soil types explain the variability of cover crop effectiveness in the temperate regions. Previous studies found the effectiveness of cover crops to reduce N leaching is highly variable, both across and within different climatic zones (Thorup-Kristensen et al., 2003; Tonitto et al., 2006; Quemada et al., 2013). In this study, soil texture had no significant impacts on N leaching under cover crops. In a review by Valkama et al. (2015) a similar relative reduction (%) in N leaching losses by cover crops, compared to the controls, across different soil textures in the Nordic countries was reported. By contrast, Premrov et al. (2014) concluded that, under mild temperate winter conditions, the risk of N leaching from light textured, freely draining soils is high and therefore, it is important to establish over-winter cover crops. In the driest parts of south-east England, early sown cover crops were found to be most effective on freely drained sandy soils, where the risk of N leaching was high, but were less effective on medium-heavy textured soils with poorer drainage (Macdonald et al., 2005).

Under cover crops, soils with higher BD are the most likely to have higher SOC. The presence of N in soil is important for SOC accumulation as C sequestration requires N (van Groenigen et al., 2017). According to Aula et al. (2016), the use of N fertilizer significantly increases SOC. The difference in SOC (%) between the cover crops and the control treatments was at its highest at low N fertilizer rate. High soil pH decreases the efficiency of cover crops to accumulate SOC. Parfitt et al. (2014) reported that high pH (due to liming) possibly reduces SOC. Both soil texture and tillage had no significant impacts on the efficiency of cover crops to sequester SOC, compared to control treatments. Previous studies showed both beneficial (West and Post, 2002; Gonzalez-Sanchez et al., 2012) and no impact (Dimassi et al., 2014; Powlson et al., 2014) of no-till relative to conventional tillage on SOC. Soil organic matter and organic residues are the two main energy sources of microbial biomass (Brookes et al., 2008). Higher SOC is advantageous for soil fertility, water holding capacity and nutrient retention and therefore, is considered essential for sustainable agriculture (Hoyle, 2013).

*Impacts of cover crops (legume, non-legume and legume-non-legume mixed) on grain yield and N content in grain of the primary crop*

We found, overall, cover crops decreased grain yields of the primary crop by  $\approx 3.9\%$  compared to the control treatment. Both legume and non-legume cover crops decreased grain yields but legume- non-legume mixed cover crops increased yield significantly. Studies found that grain yields of the primary crop can be improved by incorporation of legume-non-legume mixtures (Doltra and Olesen, 2013) or legume (Campiglia et al., 2011) cover crops. A review by Tonitto et al. (2006) reported a 10% reduction in grain yield of primary crops under legume cover crops. In contrast, Coombs et al. (2017) found alfalfa and red clover (legume) had a positive impact on corn yield in one of two years. Dozier et al. (2017) and Marcillo and Miguez (2017) found non-legume cover crops had no effects on the grain yield of corn, especially in the short

term. Noland et al. (2018) found that to reduce soil NO<sub>3</sub> while maintaining corn and subsequent soybean yields, cover crops should be inter-seeded into corn at the seven-leaf collar stage. Nevertheless, a successful termination for the cover crops is crucial to avoid competition with the subsequent soybean crop. The legume cover crop increased N in the grain of the primary crop, while non-legumes decreased it and legume-non-legume mixed cover crops had no significant effect. Wittwer et al. (2017) found higher grain N concentrations and N contents under both legume and legume-non-legume mixed cover. However, there are mixed results concerning the effects of cover crops on N content in grain of the primary crop in the literature (Thomsen, 2005; Olesen et al., 2007; Rinnofner et al., 2008; Kramberger et al., 2009; Doltra and Olesen, 2013).

#### *Impacts of cover crops (legume, non-legume and legume-non-legume mixed) on net greenhouse gas balance*

Characterising the effects of cover crops on the NGHGB of cropping systems is complex given that they influence both the carbon balance as well as direct and indirect N<sub>2</sub>O emissions. The uncertainty in our results, due to assumptions made, was conservatively estimated by calculating the standard deviations (StDev) for all values. Our study showed that all cover crop types could contribute to ecological intensification and climate change mitigation by improving the NGHGB, compared to the control treatment. Cover crop practices could also contribute to the aspirations of the soil C “4-per-mille” initiative (Minasny et al., 2017), especially in wet regions where C stocks are low and nutrients are available (e.g. North Europe, North China and Canada). The growing cover crops could increase water use, keeping soils dry and thereby reduce rates of SOC decomposition, as well as reducing N<sub>2</sub>O loss and soil erosion (Desjardins et al., 2005). In contrast, Negassa et al. (2015) reported that the addition of cover crop inputs to topographic depression areas can increase the priming effect (Guenet et al., 2010), which

increases decomposition of native SOC, and thereby increases CO<sub>2</sub> emissions, when stimulated by additions of fresh plant residue inputs. However, Steele et al. (2012) reported no changes in organic matter (OM) content after 13 years of a cover crop experiment. One limitation of our analysis is that the majority of the studies collected were short-term experiments (2-3 years). Berntsen et al. (2006) reported that the effects of cover crops should be evaluated in the long-term rather than considering short-term effects only; however, there is a scarcity of such long-term experiments. We found that incorporating cover crops, specifically legume-non-legume mixed cover crops, into the crop rotation is beneficial for soils, the environment and crop productivity. Tonitto et al. (2006) found that the legume-non-legume mixed cover crops useful for both atmospheric N<sub>2</sub> fixation and for soil residual nitrate recycling. Cover crops influence soil N and C dynamics and N available for the subsequent primary crop. They play an important role in achieving more diverse and multifunctional agricultural systems (Schipanski et al., 2014; Blanco-Canqui et al., 2015), suggesting that further efforts are required to enable farmers to overcome all barriers for their widespread adoption (Roesch-McNally et al., 2017). However, management practices in relation to cover crops will need to be adapted to specific soil, management and regional climatic conditions.

## **Concluding remarks**

This global critical review and systematic analysis reveals that, by adopting cover crops we could decrease N leaching to ground water and increase SOC sequestration without having significant effects on direct N<sub>2</sub>O emissions. To avoid the negative impacts of cover crops on grain yield (-3.9%), legume-non-legume mixed cover crops, which increase the yield by ≈13% and had no significant impacts on N in grain, should be selected. Overall, cover crops can mitigate net greenhouse gas balance by 2.06 ±2.10 Mg CO<sub>2</sub>-eq ha<sup>-1</sup> y<sup>-1</sup>. These effects can be

considered important in contributing to the resilience of farming systems to environmental changes, for example from climate change, by being more fertile, productive and have better water quality. However, to increase the effectiveness of cover crops, field management techniques should be optimized to the local climatic conditions, water resources, soil and cropping systems. The genetics of cover crop species could be improved, to provide deeper rooted crops, which have higher N use efficiencies, better nitrate scavenging abilities and lower N leaching potential. Deep rooted species could help with cover crop resilience, e.g. deeper delivery of C in the soil profile. It is also important to adjust timings and dates of the planting and kill dates of the cover crops, to avoid competition with the primary crop, to improve their effectiveness and avoid trying to establish cover crops when soil conditions are sub-optimal (potentially increasing soil erosion losses). Although cover crops increase costs, due to the need to purchase new seeds, management operations and termination costs, these costs can be compensated for if the wider benefits are considered. These include retention and carryover of nutrients between phases of a rotation, and the opportunity for the cover crops to be sold as forage or grazed. A positive return from cover crops for producers is a possibility, especially if they replace a fallow period instead of a primary crop. However, to support the widespread adoption of cover crops, improved policy, education, training and awareness raising of the potential benefits and risks and risk abatement strategies are needed.

## **Acknowledgements**

This work contributes to the UK-China Virtual Joint Centre N-Circle (grant number BB/N013484/1), SuperG (funded under EU Horizon 2020 programme)) and ADVENT (grant number NE/M019691/1). DRC was supported by the UK-China Virtual joint Centre for Agricultural Nitrogen (CINAg, BB/N013468/1) and the UK-Brazil Virtual Joint Centre to

deliver enhanced N-use efficiency via an integrated soil-plant systems approach (NUCLEUS), which are jointly supported by Newton fund *via* UK BBSRC and NERC.

## References

- Abalos, D., Deyn, G. B., Kuyper, T.W. & van Groenigen, J. W. (2014). Plant species identity surpasses species richness as a key driver of N<sub>2</sub>O emissions from grassland. *Global Change Biology*, 20, 265- 275.
- Abdalla, M., Hastings, A., Chadwick, DR., Jones, DL., Evans, CD., Jones, MB., Rees, RM. & Smith, P. (2018). Critical review of the impacts of grazing intensity on soil organic carbon storage and other soil quality indicators in extensively managed grasslands. *Agriculture Ecosystems & Environment*, 253, 62-81.
- Abdalla, M., Hastings, A., Helmy, M., Prescher, A., Osbourne, B., Lanigan, G., Forristal, D., Killi, D., Maratha, P., Williams, M., Rueangritsarakul, K., Smith, P., Nolan, P. & Jones, M.B. (2014). Assessing the combined use of reduced tillage and cover crops for mitigating greenhouse gas emissions from arable ecosystem. *Geoderma*, 223, 9-20.
- Abdalla, M., Osborne, B., Lanigan, G., Forristal, D., Williams, M., Smith, P. & Jones, M. B. (2013). Conservation tillage systems: a review of its consequences for greenhouse gas emissions. *Soil Use and Management*, 29, 199-209.
- Allingham, K.D., Cartwright, R., Donaghy, D., Conway, J.S., Goulding, K.W. & Jarvis, S.C. (2002). Nitrate leaching losses and their control in a mixed farm system in the Cotswold Hills, England. *Soil Use and Management*, 18, 421-427.
- Aronsson, H., Stenberg, M. & Ulén, B. (2011). Leaching of N, P and glyphosate from two soils after herbicide treatment and incorporation of a ryegrass catch crop. *Soil Use and*

609           *Management*, 27, 54-68.

610   Ascott, M.J., Gooddy, D., Wang, L., Stuart, M.E., Lewis, M.A., Ward, R.S. et al. (2017).

611           Global patterns of nitrate storage in the vadose zone. *Nature Communications*, 8,

612           1416.

613   Askegaard, M. & Eriksen, J. (2008). Residual effect and leaching of N and K in cropping

614           systems with clover and ryegrass catch crops on coarse sand. *Agriculture, Ecosystems*

615           *and Environment*, 123, 99-108.

616   Askegaard, M., Olesen, J.E., Rasmussen, I.A. & Kristensen, K. (2011). Nitrate leaching from

617           organic arable crop rotations is mostly determined by autumn field management.

618           *Agriculture, Ecosystems and Environment*, 142, 149-160.

619   Askegaard, M., Olesen, J.E., Rasmussen, I.A. & Kristensen, K. (2005). Nitrate leaching from

620           organic arable crop rotations: effects of location, manure and catch crop. *Soil Use and*

621           *Management*, 21, 181-188.

622   Aula, L., Macnack, N., Jeremiah, P., Mullock, J. & Raun, W. (2016). Effect of fertilizer

623           nitrogen (N) on soil organic carbon, total N, and soil pH in long-term continuous

624           winter wheat (*Triticum Aestivum* L.). *Communications in Soil Science and Plant*

625           *Analysis*, 47, issue 7.

626   Basche, A.D., Miguez, F.E., Kaspar, T.C. & Castellano, M.J. (2014). Do cover crops increase

627           or decrease nitrous oxide emissions? A meta-analysis. *Journal of Soil and Water*

628           *Conservation*, 69, 471-482.

629   Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting Linear Mixed-Effects Models

630           Using lme4. *Journal of Statistical Software*, 67(1), 1-48.

631

632   Battany, M. & Grismer, M.E. (2000). Rainfall runoff and erosion in Napa Valley vineyards:



633 effects of slope, cover and surface roughness. *Hydrological Processes*, 14, 1289-  
634 1304.

635 Benoit, M., Garnier, J., Anglade, J. & Billen, G. (2014). Nitrate leaching from organic and  
636 conventional arable crop farms in the Seine Basin (France). *Nutrient Cycling in*  
637 *Agroecosystems*, 100, 285-299.

638 Berntsen, J., Olesen, J.E., Petersen, B.M. & Hansen, E.M. (2006). Algorithms for sensor-  
639 based redistribution of nitrogen fertilizer in winter wheat. *Precision Agriculture*, 7,  
640 65-83.

641 Blanco-Canqui, H., Shaver, T.M., Lundquist, J.L., Shapiro, C.A., Elmore, R.W., Francis,  
642 C.A. & Hergert, G.W. (2015). Cover crops and ecosystem services: Insights from  
643 studies in temperate regions. *Agronomy Journal*, 107, 2449-2474.

644 Brookes, P. C., Cayuela, M. L., Contin. M., De Nobili, M., Kemmitt, S. J., & Mondini, C.  
645 (2008). The mineralization of fresh and humified soil organic matter by the soil  
646 microbial biomass. *Waste Management*, 28(4), 716-722.

647 Buchi, L., Wendling, M., Amosse, C., Necpalova, M. & Charles, R. (2018). Importance of  
648 cover crop in alleviating negative effects of reduced soil tillage and promoting soil  
649 fertility in a winter wheat cropping system. *Agriculture, Ecosystems and*  
650 *Environment*, 256, 94-104.

651 Bundy, L.G. & Andraski, T.W. (2005). Recovery of Fertilizer Nitrogen in Crop Residues and  
652 Cover Crops on an Irrigated Sandy Soil. *Soil Science Society of America Journal*, 69,  
653 640-648.

654 Campiglia, E., Mancinelli, R., Radicetti, E. & Marinari, S. (2011). Legume cover crops and  
655 mulches: Effects on nitrate leaching and nitrogen input in a pepper crop. *Nutrient*  
656 *Cycling in Agroecosystems*, 89, 399-412.

657 Cardenas, L.M., Cuttle, S.P., Crabtree, B., Hopkins, A., Shepherd, A., Scholefield, D., & Del

658 Prado, A. (2011). Cost effectiveness of nitrate leaching mitigation measures for  
659 grassland livestock systems at locations in England and Wales. *Science of the Total*  
660 *Environment*, 409 (3-4), 1104-1115.

661 Chambers, B.J., Smith, K.A. & Pain B.F. (2000). Strategies to encourage better use of  
662 nitrogen in animal manures. *Soil Use and Management*, 16, 157-166.

663 Cherr, C.M., Scholberg, J.M.S. & McSorley, R. (2006). Green manure approaches to crop  
664 production: A synthesis. *Agronomy Journal*, 98, 302-319.

665 Coombs, C., Lauzon, J.D., Deen, B. & Van Eerd, L.L. (2017). Legume cover crop  
666 management on nitrogen dynamics and yield in grain corn systems. *Field Crops*  
667 *Research*, 20, 75-85.

668 CTS. (2011). Conservation tillage service. Number 4. Available at:  
669 <http://cropsoil.psu.edu/extension/ct/uc127.pdf>; accessed on 10/01/2012.

670 Dabney, S.M., Delgado, J.A., Meisinger, J.J., Schomberg, H.H., Liebig, M.A., Kaspar, T., et  
671 al. (2011). Using cover crops and cropping systems for nitrogen management. In: J.A.  
672 Delgado and R.F. Follet, editors, Advances in nitrogen management for water quality.  
673 *Soil and Water Conservation*, Soc., Ankeny, IA. p. 230-281.

674 Defra (2010). Department for Environment Food and Rural Affairs, 2010. Fertiliser manual  
675 (RB209). <https://www.gov.uk/government/publications/fertiliser-manual-rb209--2>  
676 (accessed on 16/11/2018).

677 Delgado, J.A., Shaffer, M.J., Lal, H., McKinney, S., Gross, C.M. & Cover, H. (2008).  
678 Assessment of nitrogen losses to the environment with a Nitrogen Trading Tool  
679 (NTT). *Computer and Electronics in Agriculture*, 63, 193-206.

680 Delignette-Muller, M.L. & Dutang, C. (2015). fitdistrplus: An R package for fitting  
681 distributions. *Journal of Statistical Software*, 64(4), 1-34.

682 Desjardins, R.L., Smith, W., Grant, B., Campbell, C. & Riznek, R. (2005). Management

strategies to sequester carbon in agricultural soils and to mitigate greenhouse gas emissions. *Climatic Change*, 70, 283-297.

Dimassi, B., Mary, B., Wylleman, R., Labreuche, J., Couture, D., Piraux, F., et al. (2014). Long-term effect of contrasted tillage and crop management on soil carbon dynamics during 41 years. *Agriculture, Ecosystems and Environment*, 188, 134-46.

Ding, G., Liu, X., Herbert, S., Novak, J., Amarasiriwardena, D. & Xing, B. (2006). Effect of cover crop management on soil organic matter. *Geoderma*, 130, 229-239.

Doltra, J. & Olesen, J. (2013). The role of catch crop in the ecological intensification of spring cereals in organic farming under Nordic climate. *European Journal of Agronomy*, 44, 98-108.

Dozier, I.A., Behnke, G.D., Davis, A.S., Nafziger, E.D. & Villamil, M.B. (2017). Tillage and Cover Cropping Effects on Soil Properties and Crop Production in Illinois. *Agronomy Journal*, 109, 1261-1270.

Drinkwater, L.E. & Snapp, S.S. (2007). Nutrients in agroecosystems: Rethinking the management paradigm. *Advances in Agronomy*, 92, 63-186.

Fan, J., Hao, M. & Malhi, S.S. (2010). Accumulation of nitrate-N in the soil profile and its implications for the environment under dryland agriculture in northern China: A review. *Canadian Journal of Soil Science*, 90, 429-440.

Finney, D. M., White, C. M., & Kaye, J. P. (2016). Biomass Production and Carbon/Nitrogen Ratio Influence Ecosystem Services from Cover Crop Mixtures. *Agronomy Journal*, 108(1), 39-52.

Fraser, P.M., Curtin, D., Harrison-kirk, T., Meenken, E.D., Beare, M.H., Tabley, F., Gillespie, R.N. & Francis, G.S. (2013). Winter nitrate leaching under different tillage and winter cover crop management practices. *Soil Science Society of America Journal*, 77, 1391-1401.

708 Gonzalez-Sanchez, E.J., Ordonez-Fernandez, R., Carbonell-Bojollo, R., Veroz-Gonzalez, O.  
709 & Gil-Ribes, J.A. (2012). Meta-analysis on atmospheric carbon capture in Spain  
710 through the use of conservation agriculture. *Soil and Tillage Research*, 122, 52-60.

711 Grieser, J., Gommers, R. & Bernardi, M. (2006). The Miami Model of Climatic Net Primary  
712 Production of Biomass. The Agromet Group, SDRN, FAO of the UN, Viale delle  
713 Terme di Caracalla, 00100 Rome, Italy.

714 Guardia, G., Abalos, D., Garcia-Marco, S., Quemada, M., Alonso-Ayuso, M., Cárdenas, L.  
715 M., et al. (2016). Effect of cover crops on greenhouse gas emissions in an irrigated  
716 field under integrated soil fertility management. *Biogeosciences*, 13, 5245-5257.

717 Guenet, B., Neill, C., Bardoux, G. & Abbadie, L. (2010). Is there a linear relationship  
718 between priming effect intensity and the amount of organic matter input? *Applied Soil*  
719 *Ecology*, 46, 432-442.

720 Guo, L.B. & Gifford, R.M. (2002). Soil carbon stocks and land use change: a meta-analysis.  
721 *Global Change Biology*, 8, 345-360.

722 Halde, C., Gulden, R.H. & Entz, M.H. (2014). Selecting cover crop mulches for organic  
723 rotational no-till systems in Manitoba, Canada. *Agronomy Journal*, 106, 1193-1204.

724 Harunaa, S.I. & Nkongolo, N.V. (2015). Cover Crop Management Effects on Soil Physical  
725 and Biological Properties. *Procedia Environmental Sciences*, 29, 13-14.

726 Hontoria, C., Garcia-Gonzalez, I., Quemada, M., Roldan, A. & Alguacil, M.M. (2019).  
727 The cover crop determines the AMF community composition in soil and in roots of  
728 maize after a ten-year continuous crop rotation. *Science of The Total Environment*,  
729 660, 913-922.

730 Hooker, K.V., Coxon, C.E. Hackett, R., Kirwan, L.E., O’Keeffe, E. & Richards, K.G. (2008).  
731 Evaluation of cover crop and reduced cultivation for reducing nitrate leaching in  
732 Ireland. *Journal of Environmental Quality*, 37, 138-145.

733 IPCC (2013). Summary for policymakers. In: Stocker, T.F., Qin, D., Plattner, G.K., Tignor,  
 734 M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (eds).  
 735 Climate change 2013: the physical science basis. Contribution of Working Group I to  
 736 the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.  
 737 Cambridge University Press, Cambridge.

738 IPCC (2006). IPCC guidelines for national greenhouse gas inventories. Institute for Global  
 739 Environment Strategies, Hayama, Japan.

740 Jobbágy, E.G. & Jackson, R.B. (2001). The distribution of soil nutrients with depth: global  
 741 patterns and the imprint of plants. *Biogeochemistry*, 53, 51-77.

742 Jobbágy, E.G. & Jackson, R.B. (2000). The vertical distribution of soil organic carbon and its  
 743 relation to climate and vegetation. *Ecological Applications*, 10, 423-436.

744 Justes, E., Beaudoin, N., Bertuzzi, P., Charles, R., Constantin, J., Dürr, C., Hermon, C.,  
 745 Joannon, A., Le Bas, C., Mary, B., Mignolet, C., Montfort, F., Ruiz, L., Sarthou, J.P.,  
 746 Souchère, V., Tournebize, J., Savini, I. & Réchauchère, O. (2012). The use of cover  
 747 crops in the reduction of nitrate leaching: Impact on the water and nitrogen balance  
 748 and other ecosystem services. Summary of the study report, INRA (France), 60 pp.

749 Känkänen, H. & Eriksson, C. (2007). Effects of undersown crops on soil mineral N and grain  
 750 yield of spring barley. *European Journal of Agronomy*, 27, 25-34.

751 Känkänen, H., Eriksson, C., Räkköläinen, M. & Vuorinen, M. (2003). Soil nitrate N as  
 752 influenced by annually undersown cover crops in spring cereals. *Agricultural and*  
 753 *Food Science in Finland*, 12 (3-4), 165-176.

754 Känkänen, H., Eriksson, C., Räkköläinen, M. & Vuorinen, M. (2001). Effect of annually  
 755 repeated under-sowing on cereal grain yields. *Agricultural and Food Science in*  
 756 *Finland*, 10, 197-208.

757 Karlsson-Strese, E.M., Rydberg, I., Becker, H.C. & Umaerus, M. (1998). Strategy for catch

758 crop development II. Screening of species undersown in spring barley (*Hordeum*  
759 *vulgare* L.) with respect to catch crop growth and grain yield. *Acta Agriculturae*  
760 *Scandinavica*, 48, 26-33.

761 Kaspar, T.C., Jaynes, D.B., Parkin, T.B., Moorman, T.B. & Singer, J.W. (2012).  
762 Effectiveness of oat and rye cover crops in reducing nitrate losses in drainage water.  
763 *Agricultural Water Management*, 110, 25-33.

764 Kaspar, T.C. & Singer, J.W. (2011). The use of cover crops to manage soil. In: Hatfield, J.L.,  
765 Sauer, T.J. (Eds.), *Soil Management: Building a Stable Base for Agriculture*.  
766 American Society of Agronomy and Soil Science Society of America Journal, pp 321-  
767 337. Madison, WI.

768 Kim, Y., Seo, Y., Kraus, D., Klatt, S., Haas, E., Tenhunen, J. & Kiese, R. (2015). Estimation  
769 and mitigation of N<sub>2</sub>O emission and nitrate leaching from intensive crop cultivation in  
770 the Haeon catchment, South Korea. *Science of the Total Environment*, 529, 40-53.

771 Kramberger, B., Gselman, A., Janzekovic, M., Kaligalic, M. & Bracko, B. (2009). Effects of  
772 cover crops on soil mineral nitrogen and on the yield and nitrogen content of maize.  
773 *European Journal of Agronomy*, 31, 103-109.

774 Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. (2017). lmerTest package: tests in  
775 linear mixed effects models. *Journal of Statistical Software*, 82(13), 1-26.

776 Lee, J., Six, J., King, A.P., Van kessel, C. & Rolston, D.E. (2006). Tillage and field scale  
777 controls on greenhouse gas emissions. *Journal of Environmental Quality*, 35, 714-  
778 725.

779 Leith, H. (2000). Modelling the primary productivity of the world. *Nature and Resources*  
780 VIII. UNESCO, pp 5-10.

781 Leslie, A.W., Wang, K.H., Meyer, S.L.F., Marahatta, S. & Hooks, C.R.R. (2017). Influence

782 of cover crops on arthropods, free-living nematodes, and yield in a succeeding no-till  
 783 soybean crop. *Applied Soil Ecology*, 117-118, 21-31  
 784 Liebig, M.A., Hendrickson, J.R., Archer, D.W., Schmer, M.A., Nichols, K.A. & Tanaka,  
 785 D.L. (2015). Short-term soil responses to late-seeded cover crops in a semi-arid  
 786 environment. *Agronomy Journal*, 107, 2011-2019.  
 787 Lotter, D.W., Seidel, R. & Liebhardt, W. (2003). The performance of organic and  
 788 conventional cropping systems in an extreme climate year. *American Journal of*  
 789 *Alternative Agriculture*, 18, 146-154.  
 790 Lüscher, A., Mueller-Harvey, I., Soussana, J.F., Rees, R.M. & Peyraud, J.L. (2014). Potential  
 791 of legume-based grassland-livestock systems in Europe: a review. *Grass Forage*  
 792 *Science*, 69 (2), 206-228.  
 793 Macdonald, A.J., Poulton, P.R., How, M.T., Goulding, K.W.T. & Powlson, D.S. (2005). The  
 794 use of cover crops in cereal based cropping systems to control nitrate leaching in SE  
 795 England. *Plant and Soil*, 273, 355-373.  
 796 Madejon, E., Murillo, J.M., Moreno, F., López, M.V., Arrue, J.L., Alvaro-Fuentes, J.  
 797 & Cantero, C. (2009). Effect of long-term conservation tillage on soil biochemical  
 798 properties in Mediterranean Spanish areas. *Soil and Tillage Research*, 105, 55-62.  
 799 Marcillo, G.S. & Miguez, F.E. (2017). Corn yield response to winter cover crops: An updated  
 800 meta-analysis. *Journal of Soil and Water Conservation*, 72 (3), 226-239  
 801 Min, J., Zhao, X., Shi, W., Xing, G. & Zhu, Z. (2011). Nitrogen Balance and Loss in a  
 802 Greenhouse Vegetable System in Southeastern China. *Pedosphere*, 21 (4), 464-472.  
 803 Minasny, B., et al. (2017). Soil carbon 4 per mille. *Geoderma*, 292, 59-86.  
 804 Mitchell, D.C., Castellano, M.J., Sawyer, J.E. & Pantoja, J. (2013). Cover crop effects on  
 805 nitrous oxide emissions: Role of mineralizable carbon. *Soil Science Society of*  
 806 *America Journal*, 77, 1765-1773.

807 Negassa, W., Price, R.F., Basir, A., Snapp, S.S. & Kravchenko, A. (2015). Cover crop and  
808 tillage systems effect on soil CO<sub>2</sub> and N<sub>2</sub>O fluxes in contrasting topographic  
809 positions. *Soil and Tillage Research*, 154, 64-74.

810 Nielsen, D.C. & Vigil, M.F. (2005). Legume green fallow effect on soil water content at  
811 wheat planting and wheat yield. *Agronomy Journal*, 97, 684-689.

812 Noland, R.L., Wells, M.S., Sheaffer, C.C., Baker, J.M., Martinson, K.L. & Coulter,  
813 J.A. (2018). Establishment and Function of Cover Crops Inter-seeded into Corn. *Crop*  
814 *Science*, 58, 863-873.

815 Olson, K., Ebelhar, S.A. & Lang, J.M. (2014). Long-term effects of cover crops on crop  
816 yields, soil organic carbon stocks and sequestration. *Open Journal of Soil Science*, 4,  
817 284-292.

818 Olesen, J.E., Hansen, E.M., Askegaard, M. & Rasmussen, I.A. (2007). The value of catch  
819 crops and organic manure for spring barley in organic arable farming. *Field Crop*  
820 *Research*, 100, 168-178.

821 Parfitt, J.M.B., Timm, L.C., Reichardt, K. & Pauletto, E.A. (2014). Impacts of land levelling  
822 on lowland soil physical properties. *Revista Brasileira de Ciência do Solo*, 38, 315-  
823 326.

824 Parkin, T.B., Kaspar, T.C., Jaynes, D.B. & Moorman, T.B. (2016). Rye cover crop effects on  
825 direct and indirect nitrous oxide emissions. *Soil Science Society of America Journal*,  
826 80, 1551-1559.

827 Poeplau, C. & Don, A. (2015). Carbon sequestration in agricultural soils via cultivation of  
828 cover crops-A meta-analysis. *Agriculture, Ecosystems and Environment*, 200, 33-41.

829 Powlson, D.S., Stirling, C.M., Jat, M., Gerard, B.G., Palm, C.A., Sanchez, P.A., et al. (2014).  
830 Limited potential of no-till agriculture for climate change mitigation. *Nature Climate*  
831 *Change*, 4(8), 678-83.



832 Premrov, A., Coxon, C., Hackett, R., Kirwan, L. & Richards, K. (2014). Effects of over-  
833 winter green cover on soil solution nitrate concentrations beneath tillage land. *Science*  
834 *of the Total Environment*, 470-471, 967-974.

835 Quemada, M., Baranski, M., Nobel-de Lange, M.N.J., Vallejo, A. & Cooper, J.M. (2013).  
836 Meta-analysis of strategies to control nitrate leaching in irrigated agricultural systems  
837 and their effects on crop yield. *Agriculture, Ecosystems and Environment*, 174, 1-10.

838 Rinnofnier, T., Friedel, J.K., de Kruijff, R., Pietsch, G. & Freyer, B. (2008). Effect of catch  
839 crops on N dynamics and following crops in organic farming. *Agronomy for*  
840 *Sustainable Development*, 28, 551-558.

841 Roesch-McNally, G.E., Basche, A.D., Arbuckle, J.G., Tyndall, J.C., Miguez, F.E., Bowman,  
842 T. & Clay, R. (2017). The trouble with cover crops: Farmers' experiences with  
843 overcoming barriers to adoption. *Renewable Agriculture and Food Systems*, 1-12.

844 Schipanski, M.E., Barbercheck, M., Douglas, M.R., Finney, D.M., Haider, K., Kaye, J.P.,  
845 Kemanian, A.R., Mortensen, D.A., Ryan, M.R., Tooker, J. & White, C. (2014). A  
846 framework for evaluating ecosystem services provided by cover crops in  
847 agroecosystems. *Agricultural Systems*, 125, 12-22.

848 Smeaton, D.C., Cox, T., Kerr, S. & Dynes, R. (2011). Relationships between farm  
849 productivity, profitability, N leaching and GHG emissions: a modelling approach.  
850 *New Zealand Grasslands Association*, 73, 57-62.

851 Smith, P., Goulding, K.W.T., Smith, K.A., Powlson, D.S., Smith, J.U., Falloon, P. & Coleman,  
852 K. (2000). Including trace gas fluxes in estimates of the carbon mitigation potential of  
853 UK agricultural land. *Soil Use and Management*, 16, 251-259.

854 Smith, E. G., Peters, T.L., Blackshaw, R.E., Lindwall, C.W. & Larney, F.J. (1996).  
855 Economics of reduced tillage in crop-fallow systems. *Canadian Journal of Soil*  
856 *Science*, 76, 411-416.

857 Snapp, S.S., Swinton, S.W., Labarta, R., Mutch, D., Black, J.R., Leep, R., Nyiraneza, J. &  
858 O'Neil, K. (2005). Evaluating cover crops for benefits, costs and performance within  
859 cropping system niches. *Agronomy Journal*, 97, 322-332.

860 Steele, M.K., Coale, F.J. & Hill, R.L. (2012). Winter annual cover crop impacts on no-till soil  
861 physical properties and organic matter. *Soil Science Society of America Journal*, 76,  
862 2164-2173.

863 Steenwerth, K. & Belina, K.M. (2008). Cover crops and cultivation: Impacts on soil N  
864 dynamics and Micro-biological function in a Mediterranean vineyard agroecosystem.  
865 *Applied Soil Ecology*, 40, 370-380.

866 Stevenson, F.J. (1982). Humus Chemistry: Genesis, Composition, Reactions. Wiley-  
867 Interscience, New York.

868 Thomsen, I.K. & Hansen, E.M. (2014). Cover crop growth and impact on N leaching as  
869 affected by pre- and postharvest sowing and time of incorporation. *Soil Use and*  
870 *Management*, 30, 48-57.

871 Thomsen, I.K. (2005). Nitrate leaching under spring barley is influenced by the presence of a  
872 ryegrass catch crop: results from a lysimeter experiment. *Agriculture Ecosystems, and*  
873 *Environment*, 111, 21-29.

874 Thorup-Kristensen, K., Magid, J. & Jensen, L.S. (2003). Catch crops and green manures as  
875 biological tools in nitrogen management in temperate zone. *Advances in Agronomy*,  
876 79, 227-302.

877 Tonitto, C., David, M.B. & Drinkwater, L.E. (2006). Replacing bare fallow with cover crops  
878 in fertilizer-intensive cropping systems: a meta-analysis of crop yield and N  
879 dynamics. *Agriculture Ecosystems, and Environment*, 112, 58-72.

880 Torstensson, G. & Aronsson, H. (2000). Nitrogen leaching and crop availability in manured  
881 catch crop systems in Sweden. *Nutrient Cycling in Agroecosystems*, 56, 139-152.

882 Tribouillois, H., Constantin, J. & Justes, E. (2018). Cover crops mitigate direct greenhouse  
883 gases balance but reduce drainage under climate change scenarios in temperate  
884 climate with dry summers. *Global Change Biology*, 24 (6), 2513-2529.

885 Unger, P.W. & Vigil, M.F. (1998). Cover crop effects on soil water relationships. *Journal of*  
886 *Soil and Water Conservation*, 53, 200-207

887 Valkama, E., Lemola, R., Känkänen, H. & Turtola, E. (2015). Meta-analysis of the effects of  
888 under-sown catch crops on nitrogen leaching loss and grain yields in the Nordic  
889 countries. *Agriculture Ecosystems, and Environment*, 203, 93-101.

890 Van Groenigen, J.W, van Kessel, C., Hungate, B.A., Oenema, O., Powlson, D.S. & Van  
891 Groenigen, K.J. (2017). Sequestering Soil Organic Carbon: a Nitrogen Dilemma.  
892 *Environmental Science and Technology*, 51(9), 4738-4739.

893 Villamil, M.B., Bollero, G.A., Darmody, R.G., Simmons, F.W. & Bullock, D.G. (2006). No-  
894 till corn/soybean systems including winter cover crops. *Soil Science Society of*  
895 *America Journal*, 70, 1936.

896 Wallgren, B. & Lindén, B. (1994). Effect of catch crops and ploughing times on soil mineral  
897 nitrogen. *Swedish Journal of Agricultural Research*, 24, 67-75.

898 Wang, G. & Ngouajio, M. (2008). Integration of cover crop, conservation tillage, and low  
899 herbicide rate for machine-harvested pickling cucumbers. *HortScience*, 43(6), 1770-  
900 1774.

901 Webb, J., Harrison, R. & Ellis, S. (2000). Nitrogen fluxes in three arable soils in the UK.  
902 *European Journal of Agronomy*, 13, 207-223.

903 Weinert, T.L., Pan, W.L., Moneymaker, M.R., Santo, G.S. & Stevens, R.G. (2002). Nitrogen  
904 recycling by non-leguminous winter cover crops to reduce leaching in potato  
905 rotations. *Agronomy Journal*, 94, 365-372.

906 West, T.O. & Post, W.M. (2002). Soil organic carbon sequestration rates by tillage and crop

907 rotation. *Soil Science Society of America Journal*, 66(6), 1930-46.

908 White, C. M., Finney, D. M., Kemanian, A. R., & Kaye, J. P. (2016). A Model-Data Fusion  
 909 Approach for Predicting Cover Crop Nitrogen Supply to Corn. *Agronomy Journal*,  
 910 108, 2527-2540.

911 Wittwer, R.A., Dorn, B., Jossi, W. & van der Heijden, M.G.A. (2017). Cover crops support  
 912 ecological intensification of arable cropping systems. *Scientific Reports*, 7, 41911.

913 Wortman, S.E., Francis, C.A., Bernards, M.L., Drijber, R.A. & Lindquist, J.L. (2012).  
 914 Optimizing cover crop benefits with diverse mixtures and an alternative termination  
 915 method. *Agronomy Journal*, 104, 1425-1435.

916 Yogesh, A. & Juo, A.S.R. (1982). Leaching of fertilizer ions in a Kaolinitic Ultisol in the  
 917 high rain fall tropics: Leaching of nitrate in field plots under cropping and bare fallow.  
 918 *Soil Science Society of America Journal*, 46, 1212-1217.

919 Zhou, M. & Butterbach-Bahl, K. (2013). Assessment of nitrate leaching loss on a yield-scaled  
 920 basis from maize and wheat cropping systems. *Plant and Soil*, 374, 977-991.

921